



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

III. Solution by COOPER D. SCHMITT, A. M., Professor of Mathematics, University of Tennessee, Knoxville, Tenn.

Let the equation to tangent DE be $(x/a)\cos\phi + (y/b)\sin\phi = 1$.

Then $CB = a\sec\phi$; $CD = b\operatorname{cosec}\phi$. $DG = b + b\operatorname{cosec}\phi$.

Area $= DG \cdot GE$. $GE : CB :: DG : DC$. Whence

$$GE = \frac{a\sec\phi(1 + \operatorname{cosec}\phi)}{\operatorname{cosec}\phi} = \frac{a\sec\phi(1 + \operatorname{cosec}\phi)}{\operatorname{cosec}\phi} b(1 + \operatorname{cosec}\phi) = ab \tan\phi(1 + \operatorname{cosec}\phi)^2$$

Hence $\tan\phi(1 + \operatorname{cosec}\phi)^2$ is to be examined for a minimum. Differentiating and equating to zero, we have

$$\sec^2\phi(1 + \operatorname{cosec}\phi)^2 - 2\tan\phi(1 + \operatorname{cosec}\phi)\operatorname{cosec}\phi\cot\phi = 0,$$

whence, $\sec^2\phi(1 + \operatorname{cosec}\phi) = 2\operatorname{cosec}\phi$, or

$$\frac{1 + \sin\phi}{\sec\phi\cos^2\phi} = \frac{2}{\sin\phi}.$$

Solving, $\sin\phi = \frac{1}{2}$; whence $\operatorname{cosec}\phi = 2$.

Hence $DG = b + b\operatorname{cosec}\phi = b + 2b = 3b$.

Also solved by P. S. BERG, W. H. DRANE, J. SCHEFFER, and J. W. YOUNG.

95. Proposed by G. B. M. ZERR, A. M., Ph. D., Professor of Mathematics and Science, Chester High School, Chester, Pa.

A ship starts at the equator and sails northeast at all times. How far has the ship sailed (in miles) when her latitude is 30° , 45° , 60° , 90° ? How far when her longitude is 90° , 180° , 270° , 360° ? Regarding the earth as a sphere, radius 3956 miles.

Solution by the PROPOSER.

Let A be the point of the ship's departure, $APQR$ the ship's course, P, Q two consecutive points on the course, $AG = \theta =$ longitude of P , $PG = \phi =$ latitude of P , $OP = r =$ radius of the earth, $\angle PQN = \beta = \frac{1}{4}\pi$.

Then $PQ = ds$, $PN = EP \times \angle PEN = r\cos\phi d\theta$, $QN = rd\phi$.

$$\therefore ds^2 = r^2(\cos^2\phi d\theta^2 + d\phi^2).$$

$$\therefore ds = r(\cos^2\phi d\theta^2 + d\phi^2)^{\frac{1}{2}} \dots (1).$$

$$PN/QN = \tan\beta = \cos\phi d\theta/d\phi.$$

$$\therefore d\theta = \tan\beta d\phi/\cos\phi \dots (2).$$

$$(2) \text{ in } (1) \text{ gives } ds = \sqrt{(1 + \tan^2\beta)} d\phi = rd\phi/\cos\beta.$$


$$\therefore s = (r/\cos\beta) \int_0^\phi d\phi = r\phi/\cos\beta = r\phi\sqrt{2}.$$

When $\phi = \frac{1}{2}\pi$, $s = \frac{1}{2}\pi r\sqrt{2} = 2929.3817$ miles.

When $\phi = \frac{1}{4}\pi$, $s = \frac{1}{4}\pi r\sqrt{2} = 4394.07257$ miles.

When $\phi = \frac{3}{4}\pi$, $s = \frac{3}{4}\pi r\sqrt{2} = 5858.7634$ miles.

When $\phi = \frac{5}{4}\pi$, $s = \frac{5}{4}\pi r\sqrt{2} = 8788.14514$ miles.



From (2) $d\phi = \cos\phi d\theta / \tan\beta \dots\dots (3)$.

(3) in (1) gives

$$ds = r \cos\phi \sqrt{1 + \tan^2\beta} d\theta / \tan\beta = r \cos\phi d\theta / \sin\beta.$$

$$\text{From (2), } \theta = \tan^{-1} \frac{d\phi}{\cos\phi} = \tan^{-1} \log[\tan(\tfrac{1}{2}\pi + \tfrac{1}{2}\phi)].$$

$$\therefore e^{\theta \cot\beta} = \tan(\tfrac{1}{2}\pi + \tfrac{1}{2}\phi) = \frac{1 + \sin\phi}{\cos\phi}. \quad \therefore \cos\phi = \frac{2}{e^{\theta \cot\beta} + e^{-\theta \cot\beta}} = \frac{2}{e^{\theta} + e^{-\theta}},$$

since $\cot\beta = 1$.

$$\therefore s = 2r \int_0^{\theta} \frac{d\theta}{e^{\theta} + e^{-\theta}} = 2r \int_0^{\theta} 2(\tan^{-1}e^{\theta} - \tfrac{1}{2}\pi).$$

When $\theta = \tfrac{1}{2}\pi$, $s = 2r \int_0^{\tfrac{1}{2}\pi} 2(\tan^{-1}e^{\tfrac{1}{2}\pi} - \tfrac{1}{2}\pi) = r \int_0^{\tfrac{1}{2}\pi} 2(.3695185\pi) = 6494.764423$ miles.

When $\theta = \pi$, $s = 2r \int_0^{\pi} 2(\tan^{-1}e^{\pi} - \tfrac{1}{2}\pi) = r \int_0^{\pi} 2(.472506\pi) = 8304.902620$ miles.

When $\theta = \tfrac{3}{2}\pi$, $s = 2r \int_0^{\tfrac{3}{2}\pi} 2(\tan^{-1}e^{\tfrac{3}{2}\pi} - \tfrac{1}{2}\pi) = r \int_0^{\tfrac{3}{2}\pi} 2(.494281\pi) = 8687.626341$ miles.

When $\theta = 2\pi$, $s = 2r \int_0^{2\pi} 2(\tan^{-1}e^{2\pi} - \tfrac{1}{2}\pi) = r \int_0^{2\pi} 2(.498804\pi) = 8767.1239018$ miles.

Also solved by J. SCHEFFER. A somewhat different solution of this problem is given in Finkel's *Mathematical Solution Book*, page 344.

96. Proposed by W. H. CARTER, Vice President, and Professor of Mathematics, Centenary College, Jackson, La.

If $f(x) = \int f(x) dx$, find $f(x)$, the constant being zero.

Solution by W. F. SHAW, 1600 Sabine Street, Austin, Tex.

$$f(x) = \int f(x) dx. \quad df(x) = f(x) dx.$$

$$\frac{df(x)}{f(x)} = dx. \quad \log f(x) = x. \quad f(x) = e^x.$$

Also solved by W. H. DRANE, J. SCHEFFER, and G. B. M. ZERR.

MECHANICS.

87. Proposed by H. C. WHITAKER, M. E., Ph. D., Professor of Mathematics, Manual Training School, Philadelphia, Pa.

“He on his impious foes onward drove,
Drove them before him to the bounds
And crystal walls of Heaven; which opening wide
Rolled inward and a spacious gap disclosed
Into the wasteful deep; headlong themselves they threw
Down from the verge of Heaven.
Nine days they fell; Hell at last
Yawning received them whole and on them closed!”

Paradise Lost, Book VI.

Assuming Hell to be the center of the earth, and the only force acting on the lost spirits to be that of gravity due to the earth's attraction,—How far is Heaven?